

Causality vs. Ward identity in disordered electron systems

V. Janiš and J. Koloreň

*Institute of Physics, Academy of Sciences of the Czech Republic,
Na Slovance 2, CZ-18221 Praha 8, Czech Republic**

(Dated: February 2, 2008)

We address the problem of fulfilling consistency conditions in solutions for disordered noninteracting electrons. We prove that if we assume the existence of the diffusion pole in an electron-hole symmetric theory we cannot achieve a solution with a causal self-energy that would fully fit the Ward identity. Since the self-energy must be causal, we conclude that the Ward identity is partly violated in the diffusive transport regime of disordered electrons. We explain this violation in physical terms and discuss its consequences.

PACS numbers: 72.10.Bg, 72.15.Eb, 72.15.Qm

Quantum charge transport in random media is generally a rather complex problem. Unfortunately, even the simplest quantum tight-binding model of disordered noninteracting electrons does not provide for an exact solution. To find a complete solution of the Hamiltonian with random energies amounts to the determination of the spectrum of the Hamiltonian together with the spatial extension of the eigenfunctions. If the disorder is sufficiently small the eigenfunctions are extended waves enabling diffusion with a metallic conductivity. As shown by Anderson in his seminal paper on quantum diffusion, Ref. 1, strong randomness in the scattering potential can, however, destroy long-range tails of the extended eigenfunctions and cause electron localization. Localized eigenfunctions do not contribute to diffusion and drive the system insulating. After years of intensive studies, full understanding of Anderson localization has not yet been achieved and it still remains a big challenge in condensed matter theory.

Although there are efficient mathematical tools to handle finite random matrices,[2] it is more practical to use configurationally averaged Green functions to characterize solutions of random systems. Green functions have proved useful in particular in the thermodynamic limit where the volume and the number of particles are infinite. Exact solutions are mostly elusive and we have to resort to approximations. A mean-field theory of the Anderson model of disordered electrons, the coherent-potential approximation (CPA),[3] contains only extended states. Various extensions beyond the CPA are unable to include localized eigenstates either.[4] To take into account localized states, one has to treat and approximate one- and two-electron averaged Green functions simultaneously. The one-electron resolvent determines the spectrum of the solution, while the two-electron resolvent determines the spatial extension of the underlying eigenstates.

To arrive at a consistent solution with one- and two-electron averaged Green functions we have to comply with a number of restrictive consistency conditions reflecting analyticity and causality on the one side and conservation laws on the other. First of all we have to

warrant that the eigenenergies of the random Hamiltonian remain real. This is the case if the averaged resolvent $G(\mathbf{k}, z) = \langle \langle \mathbf{k} | [z\hat{1} - \hat{H}]^{-1} | \mathbf{k} \rangle \rangle_{av}$ has no poles in the upper and lower half-planes of complex energies z . We denote here $|\mathbf{k}\rangle$ the eigenfunctions of the non-random kinetic energy (Bloch waves), labelled by momenta $\mathbf{k}$. The analytic behavior of the averaged resolvent is expressed in the Kramers-Kronig relations between its real and imaginary parts. If we use the Dyson equation to represent the averaged resolvent via a self-energy $G(\mathbf{k}, z) = [z - \epsilon(\mathbf{k}) - \Sigma(\mathbf{k}, z)]^{-1}$ we can write the Kramers-Kronig relations for the self-energy

$$\Re\Sigma(\mathbf{k}, E+i\eta) = \Sigma_\infty + P \int_{-\infty}^{\infty} \frac{dE'}{\pi} \frac{\Im\Sigma(\mathbf{k}, E' + i\eta)}{E' - E}, \quad (1a)$$

$$\Im\Sigma(\mathbf{k}, E + i\eta) = -P \int_{-\infty}^{\infty} \frac{dE'}{\pi} \frac{\Re\Sigma(\mathbf{k}, E' + i\eta)}{E' - E}, \quad (1b)$$

where Σ_∞ is the constant part of the self-energy (its high-energy limit).

When constructing two-particle functions we have to obey besides analyticity also conservation laws expressed in Ward identities. Vollhardt and Wölfle proved the following integral Ward identity in the Anderson model of disordered electrons[5]

$$\Sigma(\mathbf{k}_+, z_+) - \Sigma(\mathbf{k}_-, z_-) = \frac{1}{N} \sum_{\mathbf{k}'} \Lambda_{\mathbf{k}\mathbf{k}'}^{eh}(z_+, z_-; \mathbf{q}) \times [G(\mathbf{k}'_+, z_+) - G(\mathbf{k}'_-, z_-)], \quad (2)$$

where Λ^{eh} stands for the irreducible vertex in the electron-hole scattering channel, $\mathbf{k}_\pm = \mathbf{k} \pm \mathbf{q}/2$, and $z_\pm = E \pm \omega \pm i\eta$, $\eta > 0$. The original proof of identity (2) did not reveal the conservation law reflected by this identity, but it was shown later that Eq. (2) is a consequence of probability conservation in the Hilbert space of Bloch waves.[6]

Identity (2) plays an important role in physics of disordered electrons, since it sets up a condition for the

existence of the diffusion pole in the density-density correlation function.[7, 8] The diffusion pole in the electron-hole correlation function is a hallmark for the diffusive character of charge transport in disordered systems.

Neither Eq. (2) nor Eq. (1) are automatically guaranteed in quantitative treatments of disordered electrons. In fact, different conditions must be fulfilled to comply with either Eq. (1) or Eq. (2). These conditions can be contradictory in specific situations and it need not be always possible to comply with both equations simultaneously. It is the aim of this Letter to materialize such a situation in models of disordered noninteracting electrons. The principal result of this Letter is the following **Assertion:** *If we assume electron-hole symmetry in the vertex functions and the existence of a diffusion pole in the low-energy limit of the electron-hole correlation function having the explicit form $2\pi n_F/[-i\omega + Dq^2]$, it is impossible to accommodate Eq. (2) for $z_+ - z_- \geq 0$ with causality of the self-energy, i. e., Eqs. (1).*

Before we proceed with a proof we first briefly discuss the assumptions in this assertion and illustrate in simple terms how a conflict between causality and the Ward identity (2) emerges.

To see how causality of the self-energy can be traced in the electron-hole irreducible vertex we reduce the complex frequency difference in Eq. (2) to a purely imaginary value, i. e. $z_+ - z_- = 2i\eta$ and set $q = 0$ to obtain a special case of the Ward identity

$$\begin{aligned} \Im \Sigma(\mathbf{k}, E + i\eta) \\ = \frac{1}{N} \sum_{\mathbf{k}'} \Lambda_{\mathbf{k}\mathbf{k}'}^{eh}(E + i\eta, E - i\eta) \Im G(\mathbf{k}', E + i\eta). \end{aligned} \quad (3)$$

We have denoted $\Lambda_{\mathbf{k}\mathbf{k}'}^{eh}(z_+, z_-) \equiv \Lambda_{\mathbf{k}\mathbf{k}'}^{eh}(z_+, z_-; \mathbf{0})$. Causality of the self-energy (negative semi-definiteness of its imaginary part) can then be controlled by positive semi-definiteness of the self-adjoint part of the electron-hole vertex appearing on the r.h.s. of Eq. (3). A causal self-energy can be determined from a given positive semi-definite electron-hole irreducible vertex. One uses the Ward identity to resolve the imaginary part of the self-energy from the electron-hole vertex and completes it with Eq. (1a) to determine the real part of the self-energy.

The self-energy calculated from the electron-hole irreducible vertex via Eqs. (3) and (1a) is causal and uniquely determined. We can warrant fulfillment of the Ward identity for imaginary energy differences, i. e., $\omega = 0$, in this construction. Behavior of the electron-hole vertex for $\omega \neq 0$ is irrelevant for the self-energy. Whether the self-energy from Eq. (3) satisfies the Ward identity also for real-energy differences remains to be checked.

We know that the mean-field solution (CPA) obeys the Ward identity for arbitrary difference of complex energies. It is due to locality of the CPA electron-hole irreducible vertex.[9] To find out whether Eq. (2) can

generally be fulfilled we hence have to analyze nonlocal electron-hole irreducible vertices.

The simplest nonlocal contributions to the electron-hole vertex are crossed diagrams. The self-energy is causal when the electron-hole irreducible vertex is a (positive) sum of products GG^* . Perturbation theory up to second order for the electron-hole vertex with this property reads

$$\Lambda_C^{eh} = \text{diagram 1} + \text{diagram 2} \quad (4)$$

where the dashed line stands for the local impurity potential $\lambda = c(\langle V^2 \rangle_{av} - \langle V \rangle_{av}^2)$ and the oriented solid lines are renormalized electron (retarded, upper line) and hole (advanced, lower line) averaged propagators. The approximate self-energy constructed from the vertex Λ_C^{eh} via Eq. (3) and Eq. (1a) is causal. It does not, however, obey the Ward identity (2) as can easily be checked by means of its proof.[8]

To comply with the Vollhardt-Wölfle identity (2) up to second order we have to supplement the simplest (maximally) crossed diagram with two further diagrams of the same order. That is, instead of Eq. (4) we have to consider the following electron-hole vertex

$$\Lambda_{WI}^{eh} = \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{diagram 4} \quad (5a)$$

We can find a formal diagrammatic solution for the self-energy corresponding to the vertex Λ_{WI}^{eh} :

$$\Sigma_{WI} = \text{diagram 5a1} + \text{diagram 5a2} \quad (5b)$$

The Ward identity holds for Σ_{WI} and Λ_{WI}^{eh} , but the self-energy is not causal. The imaginary part $\Im \Sigma_{WI}^R$ becomes positive for specific energies and disorder strengths. It hence does not fulfil the Kramers-Kronig relation (1a) and consequently it cannot be obtained from Eq. (5a) via Eq. (3). To amend this we have to add an infinite series of compensating diagrams (from the so called traveling clusters) so that the resulting self-energy representation generates a Herglotz function.[10]

The situation cannot be remedied by adding higher-order maximally crossed diagrams, i. e., diagrams with scatterings only between the electron and the hole where each impurity line crosses all other impurity potentials. With each maximally crossed diagram of order n , generating a causal contribution to the self-energy, we have to add n additional compensating non-causal diagrams.[8] The corrected irreducible vertex, compatible with the

Ward identity, generates a non-Herglotz self-energy and the sum of diagrams is not causal at any order of the expansion.

It is impossible to find a *diagrammatic* representation of a causal self-energy obeying Eq. (2) for real frequency differences and containing a sum of maximally crossed diagrams. Inconsistency between causality and the Ward identity in perturbation theory originates in these nonlocal diagrams. To prove incompatibility of causality with the Ward identity nonperturbatively we need an infinite sum of maximally crossed diagrams producing a singularity in the electron-hole irreducible vertex. This singularity, the Cooper pole, is generally an image of the diffusion pole in the full electron-hole vertex. The diffusion pole in the low-energy limit of the electron-hole correlation function was proved to have exactly the form assumed in the Assertion.[8]

Proof of the Assertion. The diffusion pole in the electron-hole correlation function is a residue of a pole in a more fundamental full two-particle vertex (Green) function $\Gamma_{\mathbf{k}\mathbf{k}'}(z_+, z_-; \mathbf{q})$. We denote Λ^{eh} and Λ^{ee} the irreducible vertices in the electron-hole and the electron-electron scattering channels characterized by antiparallel and parallel propagator lines, respectively. Let further I be a vertex irreducible in both channels. The following identity $\Lambda^{eh} + \Lambda^{ee} = I + \Gamma$ then holds. Since the vertex I is regular, the pole from the full vertex Γ must appear at least in one of the irreducible vertices. Due to the electron-hole symmetry, $\Lambda_{\mathbf{k}\mathbf{k}'}^{eh}(z_+, z_-; \mathbf{q}) = \Lambda_{\mathbf{k}\mathbf{k}'}^{ee}(z_+, z_-; -\mathbf{q} - \mathbf{k} - \mathbf{k}')$ both irreducible vertices share the same singularity.

The explicit singular low-energy asymptotics of the electron-hole irreducible vertex Λ^{eh} can be obtained from the sum of all maximally crossed diagrams. At zero temperature for $q = 0$ and $|\mathbf{k} + \mathbf{k}'| \rightarrow 0$ this sum yields

$$\Lambda_{\mathbf{k}\mathbf{k}'}^{sing}(z_+, z_-) \doteq \frac{2\pi n_F \lambda}{-i\Delta z \text{sign}(\Im \Delta z) + D(\mathbf{k} + \mathbf{k}')^2} . \quad (6)$$

We have denoted $\Delta z = z_+ - z_-$, n_F the density of states at the Fermi energy, and D the diffusion constant. This form of the low-energy behavior holds also for stronger scatterings on impurities. Multiple scatterings only renormalize characteristic parameters λ and D of the singular part of the electron-hole vertex for $\Delta z \rightarrow 0$. [7] Equation (6) is a generalization of the diffusion pole from real to complex frequency differences.

To demonstrate incompatibility of the Ward identity with causality of the self-energy we define a real-frequency function

$$\Delta W(\omega) = \frac{1}{N} \sum_{\mathbf{k}} [\Sigma^R(\mathbf{k}, E - \omega) - \Sigma^R(\mathbf{k}, E + \omega)] . \quad (7a)$$

From the Ward identity we obtain its representation via

the irreducible electron-hole vertex

$$\begin{aligned} \Delta W(\omega) = & \frac{-1}{N^2} \sum_{\mathbf{k}\mathbf{k}'} \{ 2i [\Lambda_{\mathbf{k}\mathbf{k}'}^{RA}(E + \omega, E) \\ & - \Lambda_{\mathbf{k}\mathbf{k}'}^{RA}(E - \omega, E)] \Im G^R(\mathbf{k}', E) + \Lambda_{\mathbf{k}\mathbf{k}'}^{RA}(E + \omega, E) \\ & \times [G_{\mathbf{k}'}^R(E + \omega) - G_{\mathbf{k}'}^R(E)] - \Lambda_{\mathbf{k}\mathbf{k}'}^{RA}(E - \omega, E) \\ & \times [G_{\mathbf{k}'}^R(E - \omega) - G_{\mathbf{k}'}^R(E)] \} . \quad (7b) \end{aligned}$$

We have used superscripts R, A to denote the complex half-planes of the energy arguments (upper, lower) from which the real value is approached. Analyticity of the self-energy as a function of energy E inside the energy bands induces analyticity of the function $\Delta W(\omega)$, i. e., all its derivatives are continuous functions and bounded at $\omega = 0$.

The vertex function Λ^{RA} is not, however, analytic and its nonanalyticity is transferred to $\Delta W(\omega)$. It is straightforward to evaluate $\Delta W^{sing}(\omega)$, the leading singular part of $\Delta W(\omega)$ in the limit $\omega \rightarrow 0$. We rescale the momenta $q \rightarrow x = q\sqrt{D/|\omega|}$ in Λ^{RA} and perform momentum integration in a d -dimensional space. The leading singular behavior is determined by the small-momentum asymptotics of the first term on the r.h.s. of Eq. (7b) with the singular electron-hole vertex from Eq. (6) for $\eta = 0^+$. We explicitly obtain

$$\begin{aligned} \Delta W_d^{sing}(\omega) \approx & K \lambda n_F^2 \\ & \times \begin{cases} \frac{1}{\omega} \left| \frac{\omega}{Dk_F^2} \right|^{d/2} & \text{for } d \neq 4l, \\ \frac{1}{\omega} \left| \frac{\omega}{Dk_F^2} \right|^{d/2} \ln \left| \frac{Dk_F^2}{\omega} \right| & \text{for } d = 4l, \end{cases} \quad (8) \end{aligned}$$

where K and k_F are a dimensionless constant of order one and the Fermi momentum, respectively.[11] Equation (7a), however, dictates that $\Delta W_d^{sing}(\omega) = 0$ almost everywhere inside the energy bands. We hence have a contradiction to the premise of validity of the Ward identity Eq. (2) for real frequency differences. **Q. E. D.**

We can see from Eq. (8) that the nonanalyticity in $\Delta W(\omega)$ strongly depends on the spatial dimension d . It has a square root character in odd dimensions, contains a sign function in dimensions $d = 2(2l+1)$ and a logarithm in $d = 4l$. The nonanalytic behavior is smoothened with increasing dimensionality of the system, i. e., the higher the dimension the higher is the derivative of $\Delta W(\omega)$ displaying an (infinite) jump at $\omega = 0$. It is the $[(d-2)/2]$ th derivative, where $[x]$ is the smallest integer greater than or equal to x . As expected, there is no nonanalyticity in the mean-field, $d = \infty$, limit of the electron-hole vertex and one cannot use a $1/d$ expansion around the CPA in two-particle functions.

The nonanalyticity in $\Delta W(\omega)$ demonstrates that we cannot simultaneously keep causality and the Ward identity. Causality of the self-energy (reality of eigenenergies of the random Hamiltonian) cannot be broken. We hence are forced to admit a violation of the Ward identity for

real frequency differences $z_+ - z_- = 2\omega \neq 0$ in theories with a causal self-energy and the Cooper pole in the electron-hole irreducible vertex.

Giving up the Ward identity has quantitative consequences for the diffusive behavior of disordered electrons. Most importantly, the asymptotic form of the diffusion/Cooper pole deviates from Eq. (6). This deviation in the low-energy asymptotics of the electron-hole correlation function can be explicitly manifested in high spatial dimensions. There we expect only a weak, negligible momentum dependence of the self-energy as well as of integrals of the vertex function. The Cooper singularity in the vertex function is sufficiently smoothed by momentum integration. If we introduce a parameter

$$A = -\frac{\partial}{\partial\omega} \left[\Sigma^R(\mathbf{k}, E + \omega) - \frac{1}{N} \sum_{\mathbf{k}'} \Lambda_{\mathbf{k}\mathbf{k}'}^{RA}(E + \omega, E) \times (G^R(\mathbf{k}', E + \omega) - G^A(\mathbf{k}', E)) \right]_{\omega=0} \quad (9)$$

we can represent the leading low-energy asymptotics of the zero-temperature electron-hole correlation function in high spatial dimensions as

$$\Phi_E^{RA}(\mathbf{q}, \omega) = \frac{1}{N^2} \sum_{\mathbf{k}\mathbf{k}'} G_{\mathbf{k}\mathbf{k}'}^{RA}(E + \omega, E; \mathbf{q}) \approx \frac{2\pi n_F}{-i\omega(1+A) + Dq^2}. \quad (10)$$

The parameter A increases with the disorder strength. In particular, it is the imaginary part of the frequency derivative of the electron-hole vertex, $\Im \partial \Lambda^{eh} / \partial \omega$, that drives the system toward the Anderson localization transition, where it eventually diverges. The weight of the diffusion pole $n_F/(1+A)$, expressing the number of electron states available for diffusion, decreases when approaching the localization transition as well as an effective diffusion constant $D/(1+A)$. They both vanish at the Anderson localization transition. There is hence no diffusion pole in the localized phase.

Violation of the Ward identity can be consistently explained and does not actually contradict any fundamental physical law.[12] The Ward identity holds only within the Hilbert space spanned by extended Bloch waves, the eigenstates of the nonrandom kinetic-energy operator.[8] Violation of the Ward identity then expresses the fact that we cannot construct a single Hilbert space for all configurations of the random potential. There are new, disorder-induced and configuration-dependent localized states *orthogonal* in the thermodynamic limit to Bloch (plane) waves. The statistical weight of these states is not negligible and hence the state vector of the electron in random media has a non-zero projection onto a space of localized states at any disorder strength. That is why the number of electrons in the space of extended states is not conserved when the disorder strength increases. The

weight of the localized states in the decomposition of the electron state vector increases and reaches its saturation at the Anderson localization transition, where the state vector becomes orthogonal to Bloch waves.

To conclude, we proved in this Letter that the Cooper pole, Eq. (6), in the electron-hole vertex leads in systems with disordered electrons to deviations from the Ward identity. The nonanalytic behavior due to the Cooper pole in the electron-hole vertex is not sufficiently smeared out by momentum integration in the Ward identity to produce a self-energy analytic in its energy argument. The observed deviation from the Ward identity in causal theories is explained by nonconservation of electron states available for diffusion, i.e., plane waves. Due to the existence of disorder-induced localized states, the electrons may escape from the Hilbert space of extended states into the space of localized ones, orthogonal to Bloch waves.

The research on this problem was carried out within a project AVOZ1-010-914 of the Academy of Sciences of the Czech Republic and supported in part by Grant No. 202/01/0764 of the Grant Agency of the Czech Republic. The authors thank V. Špička for discussions on quantum coherence and diffusion and VJ thanks D. Vollhardt for hospitality at the University of Augsburg and fruitful discussions on Ward identities and self-consistent theories of Anderson localization.

* Electronic address: janis@fzu.cz, kolorenc@fzu.cz

- [1] P. W. Anderson, Phys. Rev. **109**, 1492 (1958).
- [2] M. L. Mehta, *Random Matrices and Statistical Theory of Energy Levels* (Academic Press, New York 1967).
- [3] R. J. Elliot, J. A. Krumhansl, and P. L. Leath, Rev. Mod. Phys. **46**, 465 (1974).
- [4] A. Gonis, *Green Functions for Ordered and Disordered Systems* (North Holland, Amsterdam 1992).
- [5] D. Vollhardt and P. Wölffe, Phys. Rev. **B22**, 4666 (1980).
- [6] V. Janiš, Phys. Rev. **B64**, 115115 (2001).
- [7] D. Vollhardt and P. Wölffe, in *Electronic Phase Transitions*, W. Hanke and Yu. V. Kopayev (eds), (Elsevier Science Publishers B. V., Amsterdam 1992).
- [8] V. Janiš, J. Kolorenč, and V. Špička, Eur. Phys. J. **B35**, 77 (2003).
- [9] Note that the CPA does contain the diffusion pole, but the electron-hole symmetry of the vertex functions is broken in this approximation.
- [10] R. Mills and P. Ratanavararaksa, Phys. Rev. **B18**, 5291 (1978) and T. Kaplan, P. L. Leath, L. J. Gray, and H. W. Diehl, Phys. Rev. **B21**, 4230 (1980).
- [11] There is no nonanalyticity and hence no violation of the Ward identity for imaginary energy differences $\Delta z = 2i\eta$ in the electron-hole irreducible vertex.
- [12] Notice that the existing "nonperturbative" derivations of Ward identities in disordered systems apply only to single configurations of the random potential. The extension of the Ward identity to the averaged quantities is only intuitive and is based on a tacit assumption (not proved) that

we can select a single Hilbert space (spanned by Bloch waves) encompassing almost all random configurations.

See eg. R. Ramazashvili, Phys. Rev. B **66** 220503 (2002).